

Chapter 22 – STIs and AIDS

Guided Notes

22.1 – The Risks of Sexual Activity

Sexually Transmitted Infection -

Why are STIs considered a “Silent Epidemic”?

What are the reasons that account for the STI epidemic?

-
-
-

22.2 – Kinds of STIs

The three most common STIs:

The STI known as _____ is caused by a _____ that infects the urinary tract or vagina.

- In males this can lead to _____ or an inflammation of _____.
- In females this can lead to _____ or an inflammation of _____.

The most common VIRAL STI in the U.S. is caused by the _____ or HPV.

- Some forms of HPV can cause _____.
- HPV can cause _____ in women.

The most common STI caused by _____ in the U.S. is _____.

- In males this can cause _____

- In females this can cause _____

- If left untreated females may get an infection of the reproductive organs called _____ (PID).
- PID can lead to _____ in females.

Other STIs:

Hepatitis B and C are STI's that attack the _____.

- This is caused by a _____ that attacks the _____.
- Hepatitis may lead to _____ of the liver or _____, a condition where normal tissue is replaced by scar tissue.
- There is no cure

_____ is a bacterial STI that infects the urinary tract of males and females and the reproductive organs of females.

- Males usually have symptoms
- Females sometimes have symptoms
- Often times there are no recognizable symptoms
- Can be treated with antibiotics

Genital Herpes is a viral STI that causes _____

- In some people, symptoms are not noticeable
- There is no cure

Syphilis is

- Stage 1 –
- Stage 2 –
- Stage 3 –

Untreated syphilis can cause _____, _____,
and _____.